

# Rabbit IgG Heavy and Light Chain Cross-Adsorbed Antibody

Donkey Polyclonal Conjugate DyLight® 488

Antigen Affinity Purified

Catalog No. A120-208D2

Lot No. 17

---

<b>APPLICATIONS</b>	WB, IHC, ICC, Flow Cyt, IF										
<b>SPECIES REACTIVITY</b>	Rabbit. Minimum reactivity to bovine, chicken, goat, human, mouse, pig and rat										
<b>AMOUNT</b>	1 ml										
<b>CONCENTRATION</b>	0.5 mg/ml										
<b>STORAGE/SHELF LIFE</b>	2 – 8°C / 1 year from date of receipt										
<b>PHYSICAL STATE</b>	Liquid										
<b>BUFFER</b>	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.09% Sodium Azide										
<b>FLUOROPHORE/PROTEIN</b>	5										
<b>ISOTYPE</b>	IgG										
<b>ORIGIN</b>	USA										
<b>PRODUCTION PROCEDURES</b>	<p>Antiserum was cross adsorbed using bovine, chicken, goat, human, mouse, pig and rat immunosorbents to remove cross reactive antibodies. The antibody to rabbit IgG was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to DyLight® 488.</p> <p>Immunoglobulin concentration was determined using Beer's Law where 1 mg/mL IgG has an A280 of 1.4.</p> <p>By immunoelectrophoresis and ELISA this antibody reacts specifically with rabbit IgG and with light chains common to other rabbit immunoglobulins. No antibody was detected against non-immunoglobulin serum proteins. Less than 1% cross reactivity to bovine, chicken, goat, human, mouse, pig and rat IgG was detected. This antibody may cross react with IgG from other species.</p>										
<b>APPLICATIONS</b>	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:1,000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.</td></tr><tr><td>Immunohistochemistry</td><td>1:25 – 1:250</td></tr><tr><td>Immunocytochemistry</td><td>1:25 – 1:400</td></tr><tr><td>Flow Cytometry</td><td>1:50 – 1:200</td></tr><tr><td>Immunofluorescence</td><td>1:25 – 1:400</td></tr></table>	Western Blot	1:1,000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.	Immunohistochemistry	1:25 – 1:250	Immunocytochemistry	1:25 – 1:400	Flow Cytometry	1:50 – 1:200	Immunofluorescence	1:25 – 1:400
Western Blot	1:1,000 – 1:20,000. 5% non-fat dry milk in PBST or TBST is recommended for blocking and incubation of antibodies. BSA is not recommended.										
Immunohistochemistry	1:25 – 1:250										
Immunocytochemistry	1:25 – 1:400										
Flow Cytometry	1:50 – 1:200										
Immunofluorescence	1:25 – 1:400										
<b>APPLICATION NOTES</b>	<p>Not all listed applications have been specifically tested by our laboratory.</p> <p>DyLight® 488 is excited at 493 (in PBS) and emits at 518 (in PBS).</p> <p>DyLight® is a trademark of Thermo Fisher Scientific Inc. and its subsidiaries.</p>										
<b>ADDITIONAL INFO</b>	Please visit our website for additional product information.										

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer PhD Date: November 27 2022

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: [www.fortislife.com](http://www.fortislife.com)  
Orders: [orders@fortislife.com](mailto:orders@fortislife.com) • Support: [technical@fortislife.com](mailto:technical@fortislife.com)

1