

BLM Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_000048.1


Catalog No. A300-110A-T

GeneID 641


Lot No. A300-110A-T-5

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human, Mouse
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to BLM immobilized on solid support. The epitope recognized by A300-110A-T maps to a region between residue 1375 and the C-terminus (residue 1417) of human Bloom Syndrome using the numbering given in entry NP_000048.1 (GeneID 641).
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:2,000 – 1:10,000 Immunoprecipitation 2 – 10 µg/mg lysate
ADDITIONAL INFO	https://www.fortislifesciences.com/p/A300-110A-T Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: July 1, 2022



Detection of human BLM by western blot. *Samples:* Whole cell lysate (15 µg) from K-562, Jurkat, HeLa, MCF-7, and HEK293T cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-BLM antibody (A300-110A lot 5) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of mouse BLM by western blot. *Samples:* Whole cell lysate (10 µg) from CH27, BW5147.3, NIH 3T3, and CT26 cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-BLM antibody (A300-110A lot 5) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 75 seconds.



Detection of human BLM by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HEK293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-BLM antibody (A300-110A lot 5) used for IP at 6 µg per reaction. BLM was also immunoprecipitated by a previous lot of this antibody (A300-110A lot 4) and a second antibody against a different epitope of BLM (A300-120A). For blotting immunoprecipitated BLM, A300-110A was used at 0.04 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 seconds.