CAP-D3 Antibody





Rabbit Polyclonal

Antigen Affinity Purified RefSeq ID NP_056076.1

Catalog No. A300-604A-T Uniprot ID P42695 Lot No. A300-604A-T-5 GeneID 23310

APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 10 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to CAP-D3 immobilized on solid

PROCEDURES support.

The epitope recognized by A300-604A-T maps to a region between residue 1450 and the

C-terminus (residue 1498) of human Chromosome-Associated Protein D3 using the

numbering given in entry NP_056076.1 (GeneID 23310).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2.000 - 1:10.000

Immunoprecipitation $2 - 10 \mu g/mg$ lysate

ADDITIONAL INFO https://www.fortislife.com/p/A300-604A-T


Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: November 15, 2022

CAP-D3 Antibody A300-604A-T





Detection of human CAP-D3 by western blot. Samples: Whole cell lysate (50 μ g) from HEK293T, HeLa, Jurkat, MOLT-4, and CCRF-CEM cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-CAP-D3 antibody (A300-604A lot 5) used for WB at 0.1 μ g/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.

Detection of human CAP-D3 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 5% of IP loaded) from HEK293T cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-CAP-D3 antibody (A300-604A lot 5) used for IP at 6 μg per reaction. CAP-D3 was also immunoprecipitated by a previous lot of this antibody (A300-604A lot 4) and a second antibody against a different epitope of CAP-D3 (BL2943). For blotting immunoprecipitated CAP-D3, A300-604A was used at 0.04 μg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.