

CAP-G2 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID AAH43404.1

Catalog No. A300-605A-T

GeneID 54892

Lot No. A300-605A-T-4

APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 10 µl

CONCENTRATION 200 µg/ml

STORAGE/SHELF LIFE 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG

ORIGIN USA

PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to CAP-G2 immobilized on solid support.

The epitope recognized by A300-605A-T maps to a region between residue 1100 and the C-terminus (residue 1143) of human Chromosome-Associated Protein G2 using the numbering given in entry AAH43404.1 (GeneID 54892).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 – 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate


ADDITIONAL INFO <https://www.bethyl.com/product/A300-605A-T>

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD

Date: June 23, 2022



Detection of human CAP-G2 by western blot. *Samples:* Whole cell lysate (50 µg) from HeLa, HEK293T, MCF-7, K-562, and U2OS cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-CAP-G2 antibody (A300-605A lot 4) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of human CAP-G2 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-CAP-G2 antibody (A300-605A lot 4) used for IP at 6 µg per reaction. CAP-G2 was also immunoprecipitated by a previous lot of this antibody (A300-605A lot 3) and a second antibody against a different epitope of CAP-G2 (BL2946). For blotting immunoprecipitated CAP-G2, A300-605A was used at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.