DDX18 Antibody

Antigen Affinity Purified



Protein ID NP_006764.3

Catalog No. A300-635A-T GeneID 8886

Lot No. A300-635A-T-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 10 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to DDX18 immobilized on solid

PROCEDURES support.

The epitope recognized by A300–635A–T maps to a region between residues 1 and 50 of

human DEAD (Asp-Glu-Ala-Asp) Box Protein 18 using the numbering given in entry

NP_006764.3 (GeneID 8886).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 1 – 4 µg/mg lysate

ADDITIONAL INFO https://www.bethyl.com/product/A300-635A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.bethyl.com Orders: orders@fortislife.com • Support: technical@fortislife.com **DDX18 Antibody**



Detection of human DDX18 by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (5, 15 and 50 μg for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50 μg for WB) cells. Antibodies: Affinity purified rabbit anti–DDX18 antibody BL3015 (Cat. No. A300–635A) used for WB at 0.04 $\mu g/ml$ (A) and 1 $\mu g/ml$ (B) and for IP at 3 $\mu g/mg$ lysate. DDX18 was also immunoprecipitated using rabbit anti–DDX18 antibody BL3018 (Cat. No. A300–636A) at 3 $\mu g/mg$ lysate. Detection: Chemiluminescence with exposure times of 3 minutes (A and B).