Phospho MCM2 (S53) Antibody





Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_004517.2

Catalog No. A300-756A-T GeneID 4171

Lot No. A300-756A-T-4

APPLICATIONS WB, IHC

SPECIES REACTIVITY Human, Mouse

AMOUNT 10 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to Phospho MCM2-S53 immobilized

PROCEDURES on solid support.

Immunogen for A300-756A-T was a synthetic phosphorylated peptide, which represented a

portion of human minichromosomal maintenance deficient 2 surrounding Serine 53

according to the numbering given in entry NP_004517.2 (GeneID 4171).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:1,000 - 1:2,500

Immunohistochemistry 1:100 - 1:500. Epitope retrieval with citrate buffer pH 6.0 is

recommended for FFPE tissue sections.


ADDITIONAL INFO https://www.bethyl.com/product/A300-756A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022



Detection of human Phospho MCM2 (S53) by western blot. Samples: Whole cell lysate (5 and 15 μg for WB; 1 mg for IP, 20% of IP loaded) from asynchronous HEK293T cells. For IP/WB, MCM2 was immunoprecipitated using A300–191A, which recognizes total MCM2. The immunoprecipitate was mock treated (–) or treated (+) with phosphatase (PPase). Antibody: Affinity purified rabbit anti–phospho MCM2 (S53) antibody A300–756A (lot A300–756A–4) used at 0.1 μg/ml. To examine total MCM2, the membrane was stripped and blotted with A300–191A at 0.1 μg/ml. Detection: Chemiluminescence with exposure times of 10 seconds (A and B).



Detection of human and mouse Phospho MCM2 (S53) by immunohistochemistry. Sample: FFPE sections of human breast carcinoma (left) and mouse renal cell carcinoma (right). Antibody: Affinity purified rabbit anti-phospho MCM2 (S53) (Cat. No. A300-756A Lot4) used at a dilution of 1:200 (1µg/ml). Detection: DAB