## **FAK Antibody**

Rabbit Polyclonal



Catalog No. A301-693A-T GenelD 5747

Lot No. A301-693A-T-1

**APPLICATIONS** WB, IP SPECIES REACTIVITY Human

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Mouse and Rat

**AMOUNT** 10 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION PROCEDURES

Antibody was affinity purified using an epitope specific to FAK immobilized on solid support.

The epitope recognized by A301-693A-T maps to a region between residue 1002 and 1052 of human focal adhesion kinase 1 using the numbering given in entry NP\_005598.3 (GeneID

5747).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 – 5 μg/mg lysate

ADDITIONAL INFO https://www.bethyl.com/product/A301-693A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.bethyl.com Orders: orders@fortislife.com • Support: technical@fortislife.com FAK Antibody A301-693A-T



## Detection of human FAK by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (5, 15 and 50 $\mu$ g for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50 $\mu$ g) cells. Antibodies: Affinity purified rabbit anti-FAK antibody A301-693A used for WB at 0.04 $\mu$ g/ml (A) and 1 $\mu$ g/ml (B) and used for IP at 3 $\mu$ g/mg lysate. FAK was also immunoprecipitated by rabbit anti-FAK antibody A301-692A, which recognizes an upstream epitope. Detection: Chemiluminescence with exposure times of 30 seconds (A) and 3 seconds (B).