

BORG2 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_006440.2

Catalog No. A301-723A-T

GeneID 10602

Lot No. A301-723A-T-1

| | |
|------------------------------|---|
| APPLICATIONS | IP |
| SPECIES REACTIVITY | Human |
| PRESUMED REACTIVITY | Based on 100% sequence identity, this antibody is predicted to react with Mouse |
| AMOUNT | 10 µl |
| CONCENTRATION | 1000 µg/ml |
| STORAGE/SHELF LIFE | 2 – 8°C / 1 year from date of receipt |
| PHYSICAL STATE | Liquid |
| BUFFER | Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide |
| ISOTYPE | IgG |
| ORIGIN | USA |
| PRODUCTION PROCEDURES | Antibody was affinity purified using an epitope specific to BORG2 immobilized on solid support. |

The epitope recognized by A301-723A-T maps to a region between residue 204 and 254 of human binder of Rho GTPases 2 using the numbering given in entry NP_006440.2 (GeneID 10602).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

APPLICATIONS

Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

| | |
|--------------|--|
| Western Blot | Not recommended. A301-723A has not performed satisfactorily when used for WB of BORG2 in crude preparations (e.g. whole cell lysate). This antibody can be used for WB of enriched (e.g. immunoprecipitated) sources of BORG2. |
|--------------|--|

| | |
|---------------------|--------------------|
| Immunoprecipitation | 2 – 5 µg/mg lysate |
|---------------------|--------------------|


ADDITIONAL INFO

<https://www.bethyl.com/product/A301-723A-T>

Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human BORG2 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-BORG2 antibody A301-723A used for IP at 3 µg/mg lysate. BORG2 was also immunoprecipitated by rabbit anti-BORG2 antibody A301-722A, which recognizes a downstream epitope. For blotting immunoprecipitated BORG2, A301-723A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.