

# NUP96 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP\_057404.2

Catalog No. A301-785A-T

GeneID 4928

Lot No. A301-785A-T-1

<b>APPLICATIONS</b>	IP
<b>SPECIES REACTIVITY</b>	Human
<b>PRESUMED REACTIVITY</b>	Based on 100% sequence identity, this antibody is predicted to react with Mouse and Rat
<b>AMOUNT</b>	10 µl
<b>CONCENTRATION</b>	1000 µg/ml
<b>STORAGE/SHELF LIFE</b>	2 – 8°C / 1 year from date of receipt
<b>PHYSICAL STATE</b>	Liquid
<b>BUFFER</b>	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
<b>ISOTYPE</b>	IgG
<b>ORIGIN</b>	USA
<b>PRODUCTION PROCEDURES</b>	Antibody was affinity purified using an epitope specific to NUP96 immobilized on solid support.

The epitope recognized by A301-785A-T maps to a region between residue 1650 and 1700 of human nucleoporin 96kDa using the numbering given in entry NP\_057404.2 (GeneID 4928).

Immunoglobulin concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot Not recommended. Use rabbit anti-NUP96 antibody A301-784A.

Immunoprecipitation 2 – 5 µg/mg lysate

**ADDITIONAL INFO** <https://www.bethyl.com/product/A301-785A-T>

Use the link above to view SDS, a current list of citations, and other product specific information.  
IP-western blot protocol: [https://www.bethyl.com/content/protocol\\_IP\\_WB](https://www.bethyl.com/content/protocol_IP_WB)

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.  
Michael Spencer, PhD Date: June 6, 2022



**Detection of human NUP96 by western blot of immunoprecipitates.** *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibody:* Affinity purified rabbit anti-NUP96 antibody A301-785A used for IP at 3 µg/mg lysate. NUP96 was also immunoprecipitated by rabbit anti-NUP96 antibody A301-784A, which recognizes an upstream epitope. For blotting immunoprecipitated NUP96, A301-784A was used at 0.4 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.