

BORG5 Antibody

Rabbit Polyclonal

Antigen Affinity Purified

Protein ID NP_689449.1

Catalog No. A302-381A

GeneID 11135


Lot No. A302-381A-1



| | |
|------------------------------|--|
| APPLICATIONS | IP |
| SPECIES REACTIVITY | Human |
| AMOUNT | 100 µl |
| CONCENTRATION | 1000 µg/ml |
| STORAGE/SHELF LIFE | 2 – 8° C / 1 year from date of receipt |
| PHYSICAL STATE | Liquid |
| BUFFER | Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide |
| ISOTYPE | IgG |
| ORIGIN | USA |
| PRODUCTION PROCEDURES | <p>Antibody was affinity purified using an epitope specific to BORG5 immobilized on solid support.</p> <p>The epitope recognized by A302-381A maps to a region between residue 150 and 200 of human binder of Rho GTPases 5 using the numbering given in entry NP_689449.1 (GeneID 11135).</p> <p>Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of 1.4 equals 1.0 mg of IgG.</p> |
| APPLICATIONS | <p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <p>Western Blot Not recommended. Use rabbit anti-BORG5 antibody A302-382A.</p> <p>Immunoprecipitation 5 – 10 µg/mg lysate</p> |
| APPLICATION NOTES | Western blot of immunoprecipitates performed using Normal Pig Serum (Cat. No. S100-020), Goat anti-Rabbit Light Chain HRP Conjugate (Cat. No. A120-113P) and 4-20% SDS-PAGE (link to IP-western blot protocol in Additional Info section below). |
| ADDITIONAL INFO | <p>https://www.bethyl.com/product/A302-381A</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p> <p>IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB</p> |

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Eric McIntush, PhD | Chief Scientific Officer

Date: June 21, 2019



Detection of human BORG5 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP, 20% of IP loaded) from HeLa cells. *Antibodies:* Affinity purified rabbit anti-BORG5 antibody A302-381A used for IP at 10 µg/mg lysate. BORG5 was also immunoprecipitated by rabbit anti-BORG5 antibody A302-382A, which recognizes a downstream epitope. For blotting immunoprecipitated BORG5, A302-381A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 3 seconds.