## **NUP133 Antibody**



Antigen Affinity Purified Protein ID NP\_060700.2

Catalog No. A302-386A-T GeneID 55746

Lot No. A302-386A-T-1

APPLICATIONS WB, IP

SPECIES REACTIVITY Human

AMOUNT 10 μl

CONCENTRATION 200 μg/ml

STORAGE/SHELF LIFE 2 - 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to NUP133 immobilized on solid

**PROCEDURES** support.

The epitope recognized by A302–386A–T maps to a region between residue 1106 to 1156 of human nucleoporin 133kDa using the numbering given in entry NP\_060700.2 (GenelD

55746).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2.000 - 1:10.000

Immunoprecipitation 5 – 10 µg/mg lysate

ADDITIONAL INFO https://www.bethyl.com/product/A302-386A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.bethyl.com Orders: orders@fortislife.com • Support: technical@fortislife.com



Detection of human NUP133 by western blot and immunoprecipitation. Samples: Whole cell lysate from HeLa (5, 15 and 50  $\mu g$  for WB; 1 mg for IP, 20% of IP loaded) and HEK293T (T; 50  $\mu g$ ) cells. Antibodies: Affinity purified rabbit anti–NUP133 antibody A302–386A used for WB at 0.04  $\mu g/ml$  (A) and 1  $\mu g/ml$  (B) and used for IP at 10  $\mu g/mg$  lysate. NUP133 was also immunoprecipitated by rabbit anti–NUP133 antibody A302–385A, which recognizes an upstream epitope. Detection: Chemiluminescence with exposure times of 3 minutes (A) and 10 seconds (B).