## **MNT Antibody**

Antigen Affinity Purified



Protein ID NP 064706.1

Catalog No. A303-626A-T GeneID 4335

Lot No. A303-626A-T-2

**APPLICATIONS** WB, IP SPECIES REACTIVITY Human

PRESUMED REACTIVITY Based on 100% sequence identity, this antibody is predicted to react with Mouse

AMOUNT 10 μl

CONCENTRATION 1000 μg/ml

**STORAGE/SHELF LIFE** 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

**PRODUCTION** Antibody was affinity purified using an epitope specific to MNT immobilized on solid

**PROCEDURES** support.

The epitope recognized by A303-626A-T maps to a region between residue 1 and 50 of human MAX Binding Protein using the numbering given in entry NP\_064706.1 (GenelD

4335).

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 - 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate

ADDITIONAL INFO https://www.bethyl.com/product/A303-626A-T

Use the link above to view SDS, a current list of citations, and other product specific information.


IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.bethyl.com Orders: orders@fortislife.com • Support: technical@fortislife.com MNT Antibody A303-626A-T





Detection of human MNT by western blot. Samples: Whole cell lysate (10 μg) from HEK293T, HeLa, Hep-G2, LNCaP, and A-549 cells prepared using NETN lysis buffer. Antibody: Affinity purified rabbit anti-MNT antibody (A303-626A lot 2) used for WB at 0.1 μg/ml. Detection: Chemiluminescence with an exposure time of 3 minutes.

Detection of human MNT by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from Hep–G2 cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti–MNT antibody (A303–626A lot 2) used for IP at 6 μg per reaction. MNT was also immunoprecipitated by a previous lot of this antibody (A303–626A lot 1) and another antibody against a different epitope of MNT (A303–627A) . For blotting immunoprecipitated MNT, A303–626A was used at 0.1 μg/ml. *Detection:* Chemiluminescence with an exposure time of 1 second.