

Gamma-Taxilin Antibody

Rabbit Polyclonal

Antigen Affinity Purified	Protein ID	NP_060830.2
Catalog No. A304-042A-T	GeneID	55787
Lot No. A304-042A-T-1		

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA
PRODUCTION PROCEDURES	Antibody was affinity purified using an epitope specific to Gamma-Taxilin immobilized on solid support.

The epitope recognized by A304-042A-T maps to a region between residue 475 to 525 of human Taxilin Gamma using the numbering given in entry NP_060830.2 (GeneID 55787). The epitopes for anti-ABT1 antibodies A304-044A and A304-045A are in close proximity, but do not overlap.

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot 1:2,000 – 1:10,000


Immunoprecipitation 2 – 10 µg/mg lysate

ADDITIONAL INFO <https://www.bethyl.com/product/A304-042A-T>


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human Gamma-Taxilin by western blot.
Samples: Whole cell lysate (50 µg) from HEK293T, HeLa, and Jurkat cells. *Antibodies:* Affinity purified rabbit anti-Gamma-Taxilin antibody A304-042A (lot A304-042A-1) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of human Gamma-Taxilin by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1 mg for IP; 20% of IP loaded) from HEK293T cells. *Antibodies:* Affinity purified rabbit anti-Gamma-Taxilin antibody A304-042A (lot A304-042A-1) used for IP at 6 µg/mg lysate. Gamma-Taxilin was also immunoprecipitated by rabbit anti-Gamma-Taxilin antibodies A304-041A and A304-043A. For blotting immunoprecipitated Gamma-Taxilin, A304-042A was used at 1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.