

gamma Adaptin/AP1G1 Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID O43747

Catalog No. A304-771A-T GeneID 164

Lot No. A304-771A-T-1

APPLICATIONS	WB, IP
SPECIES REACTIVITY	Human, Mouse
PRESUMED REACTIVITY	Based on 100% sequence identity, this antibody is predicted to react with Orangutan
AMOUNT	10 µl
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide
ISOTYPE	IgG
ORIGIN	USA

PRODUCTION PROCEDURES Antibody was affinity purified using an epitope specific to gamma Adaptin/AP1G1 immobilized on solid support.

The epitope recognized by A304-771A-T maps to a region between residue 1 to 50 of human AP-1 complex subunit gamma-1 using the numbering given in entry O43747 (GeneID 164).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.


Western Blot 1:2,000 – 1:10,000

Immunoprecipitation 2 – 10 µg/mg lysate


ADDITIONAL INFO <https://www.bethyl.com/product/A304-771A-T>

Use the link above to view SDS, a current list of citations, and other product specific information. IP-western blot protocol: https://www.bethyl.com/content/protocol_IP_WB


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: June 6, 2022



Detection of human gamma Adaptin/AP1G1 by western blot. *Samples:* Whole cell lysate (50 µg) from HeLa, HEK293T, and Jurkat cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-gamma Adaptin/AP1G1 antibody A304-771A (lot A304-771A-1) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.



Detection of mouse gamma Adaptin/AP1G1 by western blot. *Samples:* Whole cell lysate (50 µg) from TCMK-1 and NIH 3T3 cells prepared using NETN lysis buffer. *Antibody:* Affinity purified rabbit anti-gamma Adaptin/AP1G1 antibody A304-771A (lot A304-771A-1) used for WB at 0.1 µg/ml. *Detection:* Chemiluminescence with an exposure time of 30 seconds.



Detection of human gamma Adaptin/AP1G1 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. *Antibodies:* Affinity purified rabbit anti-gamma Adaptin/AP1G1 antibody A304-771A (lot A304-771A-1) used for IP at 6 µg per reaction. gamma Adaptin/AP1G1 was also immunoprecipitated by rabbit anti-gamma Adaptin/AP1G1 antibody BL18432. For blotting immunoprecipitated gamma Adaptin/AP1G1, A304-771A was used at 0.4 µg/ml. *Detection:* Chemiluminescence with an exposure time of 10 seconds.