## SLX4IP/C20orf94 Antibody





Rabbit Polyclonal

Antigen Affinity Purified Protein ID Q5VYV7.1 Catalog No. A304-995A-T GeneID 128710

Lot No. A304-995A-T-1

APPLICATIONS IP

SPECIES REACTIVITY Human

AMOUNT 10 μl

CONCENTRATION 1000 μg/ml

**STORAGE/SHELF LIFE** 2 – 8°C / 1 year from date of receipt

PHYSICAL STATE Liquid

**BUFFER** Tris-citrate/phosphate buffer, pH 7 to 8 containing 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTION Antibody was affinity purified using an epitope specific to SLX4IP/C20orf94 immobilized on

**PROCEDURES** solid support.

The epitope recognized by A304-995A-T maps to a region between residue 175 to 225 of human UPF0492 protein C20orf94 using the numbering given in entry Q5VYV7.1 (GeneID

128710).

Antibody concentration was determined by extinction coefficient: absorbance at 280 nm of

1.4 equals 1.0 mg of IgG.

**APPLICATIONS** Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Western Blot Not recommended. Use rabbit anti-SLX4IP/C20orf94 antibody A304-

994A

Immunoprecipitation  $2 - 10 \mu g/mg$  lysate

ADDITIONAL INFO https://www.bethyl.com/product/A304-995A-T


Use the link above to view SDS, a current list of citations, and other product specific information.

IP-western blot protocol: https://www.bethyl.com/content/protocol\_IP\_WB

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.

Michael Spencer, PhD

Date: June 6, 2022



Detection of human SLX4IP/C20orf94 by western blot of immunoprecipitates. Samples: Whole cell lysate (0.5 or 1.0 mg per IP reaction; 20% of IP loaded) from HeLa cells prepared using NETN lysis buffer. Antibodies: Affinity purified rabbit anti-SLX4IP/C20orf94 antibody A304-995A (lot A304-995A-1) used for IP at 6 μg per reaction. SLX4IP/C20orf94 was also immunoprecipitated by rabbit anti-SLX4IP/C20orf94 antibody A304-994A. For blotting immunoprecipitated SLX4IP/C20orf94, A304-994A was used at 1 μg/ml. Detection: Chemiluminescence with an exposure time of 30 seconds.