

B7-H3/CD276 Recombinant Monoclonal Antibody [BLR025F]

Rabbit Recombinant Monoclonal

Purified	RefSeq ID	NP_001019907.1
Catalog No. A700-025	Uniprot ID	Q5ZPR3
Lot No. 3	GeneID	80381

APPLICATIONS	WB, IP, IHC, ICC, IHC-IF, Flow Cyt												
SPECIES REACTIVITY	Human, Mouse												
AMOUNT	100 µl (50+ tests)												
CONCENTRATION	1000 µg/ml												
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt												
PHYSICAL STATE	Liquid												
BUFFER	Phosphate Buffered Saline (PBS) with 0.09% Sodium Azide, BSA-Free												
ISOTYPE	IgG												
CLONE #	BLR025F												
ORIGIN	USA												
PRODUCTION PROCEDURES	<p>Recombinant antibody was purified from cell culture supernatant.</p> <p>Immunogen was a peptide representing a region between residue 500 and the C-terminus (residue 534) of human CD276 Molecule using the numbering given in entry NP_001019907.1 (GeneID 80381).</p>												
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table><tr><td>Western Blot</td><td>1:1000</td></tr><tr><td>Immunoprecipitation</td><td>20 µl/mg lysate</td></tr><tr><td>Immunohistochemistry</td><td>1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.</td></tr><tr><td>Immunocytochemistry</td><td>1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE cell sections.</td></tr><tr><td>Immunofluorescence (IHC)</td><td>1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.</td></tr><tr><td>Flow Cytometry</td><td>Fixed in 4% formaldehyde and permeabilized with 90% methanol. 1 µl per 1 x 10⁶ cells.</td></tr></table>	Western Blot	1:1000	Immunoprecipitation	20 µl/mg lysate	Immunohistochemistry	1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.	Immunocytochemistry	1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE cell sections.	Immunofluorescence (IHC)	1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.	Flow Cytometry	Fixed in 4% formaldehyde and permeabilized with 90% methanol. 1 µl per 1 x 10 ⁶ cells.
Western Blot	1:1000												
Immunoprecipitation	20 µl/mg lysate												
Immunohistochemistry	1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.												
Immunocytochemistry	1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE cell sections.												
Immunofluorescence (IHC)	1:100 – 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections.												
Flow Cytometry	Fixed in 4% formaldehyde and permeabilized with 90% methanol. 1 µl per 1 x 10 ⁶ cells.												
APPLICATION NOTES	<p>All western blot analysis is performed using 5% Milk-TBST for blocking and as antibody diluent. Primary antibody is incubated overnight.</p> <p>Western blots of cell lysates are performed using Goat anti-Rabbit IgG Heavy and Light Chain Antibody (A120-101P).</p> <p>Western blots of immunoprecipitates are performed using Goat anti-Rabbit Light Chain HRP Conjugate (A120-113P) with 5% Normal Pig Serum (S100-020) added to the blocking buffer.</p>												
IHC HUMAN CONTROLS	Breast Carcinoma, Lung Carcinoma, Ovarian Carcinoma, Prostate Carcinoma, A-172 Cells, HEK293T Cells, HeLa Cells, MCF-7 Cells												
ADDITIONAL INFO	<p>https://www.fortislife.com/p/A700-025</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>												

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: March 27, 2025

Phone: 800.338.9579 • Fax: 866.597.6105 • Web: www.fortislife.com
Orders: orders@fortislife.com • Support: technical@fortislife.com

For in vitro laboratory use only. Not for any clinical, therapeutic or diagnostic use in humans or animals. Not for human or animal consumption. This product may not be resold or modified for resale without the prior written approval of Bethyl Laboratories, Inc. The information provided in this data sheet is believed to be correct but does not purport to be all-inclusive and is intended to be used as a guide. Bethyl Laboratories, Inc. shall not be liable or responsible in any way for use of either this information or the material supplied. Disposal of hazardous material may be subject to federal, state or local laws or regulations.