


FOXP3 Recombinant Monoclonal Antibody [BLR034F]

Rabbit Recombinant Monoclonal


Purified Protein ID NP_054728.2
Catalog No. A700-034-T GenelD 50943
Lot No. A700-034-T-2

APPLICATIONS	WB, IP, IHC, ICC, IHC-IF, mIF
SPECIES REACTIVITY	Human
AMOUNT	10 µl (5+ tests)
CONCENTRATION	1000 µg/ml
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt
PHYSICAL STATE	Liquid
BUFFER	Borate Buffered Saline (BBS) pH 8.2 with 0.09% Sodium Azide, BSA Free
ISOTYPE	IgG
CLONE #	BLR034F
ORIGIN	USA
PRODUCTION PROCEDURES	Recombinant antibody was purified from cell culture supernatant. Immunogen was a peptide representing a region between residue 275 and 325 of human Forkhead box P3 using the numbering given in entry NP_054728.2 (Gene ID 50943).
APPLICATIONS	Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use. Western Blot 1:1000 Immunoprecipitation 20 ul/mg lysate Immunohistochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections. Immunocytochemistry 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE cell sections. Immunofluorescence (IHC) 1:100 to 1:500. Epitope retrieval with citrate buffer pH6.0 is recommended for FFPE tissue sections. Multiplex Immunofluorescence 1:250
IHC HUMAN CONTROLS	Breast Carcinoma, Lung Carcinoma, Prostate Carcinoma, Renal Cell Carcinoma, Tonsil, MJ Cells
ADDITIONAL INFO	https://www.fortislife.com/p/A700-034-T Use the link above to view SDS, a current list of citations, and other product specific information.


This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: August 5, 2022




Detection of human FOXP3 by western blot. *Samples:* Whole cell lysate (50 µg) from Jurkat, HEK293T, MJ, MOLT-4, and RPMI-8226 cells prepared using NETN lysis buffer. *Antibody:* Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034-T lot 2) used at 1:1000. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-101P). Chemiluminescence with an exposure time of 30 seconds. Lower Panel: Rabbit anti-COPB2 antibody (A304-523A).




Detection of human FOXP3 by western blot of immunoprecipitates. *Samples:* Whole cell lysate (1.0 mg per IP reaction; 20% of IP loaded) from MJ cells prepared using NETN lysis buffer. *Antibodies:* Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034-T lot 2) used for IP at 20 µl/mg lysate. FOXP3 was also immunoprecipitated by a previous lot of this antibody (A700-034 lot 1). For blotting immunoprecipitated FOXP3, A700-034-T was used at 1:1000. Chemiluminescence with an exposure time of 30 seconds.



Detection of human FOXP3 in FFPE tonsil by immunohistochemistry. *Antibody:* Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034-T lot 2). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P). *Substrate:* DAB.



Detection of human FOXP3 in FFPE breast carcinoma by immunohistochemistry. *Antibody:* Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034-T lot 2). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P). *Substrate:* DAB.



Detection of human FOXP3 in FFPE MJ cells by immunocytochemistry. *Antibody:* Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034-T lot 2). *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P). *Substrate:* DAB.

Detection of human FOXP3 (red) by immunohistochemistry. *Sample:* FFPE section of human breast carcinoma. *Antibody:* Rabbit anti-FOXP3 recombinant monoclonal antibody [BLR034F] (A700-034-T lot 1) used at 1:250. *Secondary:* HRP-conjugated goat anti-rabbit IgG (A120-501P). *Substrate:* Opal™. *Counterstain:* DAPI (blue).