


BETHYL | **FORTIS**
LABORATORIES, INC. | LIFE SCIENCES

APPLICATIONS	WB, IHC, ICC, ELISA								
SPECIES REACTIVITY	Human								
AMOUNT	1 ml								
CONCENTRATION	1 mg/ml								
STORAGE/SHELF LIFE	2 – 8°C / 1 year from date of receipt								
PHYSICAL STATE	Liquid								
BUFFER	Phosphate Buffered Saline (PBS) containing 0.2% BSA and 0.05% Pro-Clean 400								
ISOTYPE	IgG								
ORIGIN	USA								
PRODUCTION PROCEDURES	<p>Antiserum was solid phase adsorbed to ensure class specificity. The antibody was isolated by affinity chromatography using antigen coupled to agarose beads and conjugated to horseradish peroxidase (HRP).</p> <p>Prior to conjugation, immunoglobulin concentration was determined using Beer's Law where 1mg/mL IgG has an A280 of 1.4. Molar enzyme/antibody protein ratio is 4:1.</p> <p>By immunoelectrophoresis and ELISA this antibody reacts specifically with human IgG. Cross reactivity with IgM, IgA and light chains is less than 1%. This antibody may cross react with IgG from other species.</p>								
APPLICATIONS	<p>Centrifuge tube to remove product from lid. Optimal working dilutions should be determined experimentally by the investigator. Prepare working dilution immediately before use.</p> <table border="0"> <tr> <td>Western Blot</td> <td>1:5000 – 1:50,000</td> </tr> <tr> <td>Immunohistochemistry</td> <td>1:200 – 1:5000</td> </tr> <tr> <td>Immunocytochemistry</td> <td>1:200 – 1:5,000</td> </tr> <tr> <td>ELISA</td> <td>1:10,000 – 1:100,000</td> </tr> </table>	Western Blot	1:5000 – 1:50,000	Immunohistochemistry	1:200 – 1:5000	Immunocytochemistry	1:200 – 1:5,000	ELISA	1:10,000 – 1:100,000
Western Blot	1:5000 – 1:50,000								
Immunohistochemistry	1:200 – 1:5000								
Immunocytochemistry	1:200 – 1:5,000								
ELISA	1:10,000 – 1:100,000								
APPLICATION NOTES	Not all listed applications have been specifically tested by our laboratory.								
ADDITIONAL INFO	<p>https://www.fortislife.com/p/A80-104P</p> <p>Use the link above to view SDS, a current list of citations, and other product specific information.</p>								

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc.
Michael Spencer, PhD Date: February 14, 2024

**Detection of human IgG-Fc Fragment by western blot.**

Samples: Whole cell lysate (10 µg) from HEK293T, VLN3G2, and Jurkat cells prepared using NETN lysis buffer. *Antibody:* HRP-conjugated Goat anti-Human IgG-Fc Fragment Antibody (A80-104P) used for WB at 1 µg/ml. *Secondary:* None. *Detection:* Chemiluminescence with an exposure time of 30 seconds.