Phospho MCM2 (S108) IHC Antibody

Rabbit Polyclonal

Antigen Affinity Purified Protein ID NP_004517.2

Catalog No. IHC-00014-T GeneID 4171

Lot No. IHC-00014-T-3

APPLICATIONS IHC, IHC–IF

SPECIES REACTIVITY Human, Mouse

AMOUNT 10 μl

CONCENTRATION 100 μg/ml

STORAGE/SHELF LIFE 2 - 8° C / 1 year from date of receipt

PHYSICAL STATE Liquid

BUFFER Tris-buffered Saline containing 0.1% BSA and 0.09% Sodium Azide

ISOTYPE IgG
ORIGIN USA

PRODUCTIONAntibody was affinity purified using an epitope specific to Phospho–MCM2 (Ser 108) immobilized on solid support. Immunogen for IHC–00014–T was a synthetic phosphorylated peptide, which

on solid support. Immunogen for IHC-00014-T was a synthetic phosphorylated peptide, which represented a portion of human Minichromosomal Maintenance Deficient 2 surrounding Serine

108 according to the numbering given in entry NP_0045117.2 (GenelD 4171).

APPLICATIONS Centrifuge tube to remove product from lid. Optimal working dilutions should be determined

experimentally by the investigator. Prepare working dilution immediately before use.

Immunohistochemistry 1:100 - 1:500

Immunofluorescence 1:50 - 1:500

(IHC)

APPLICATION NOTES Epitope exposure is recommended.

Epitope exposure with citrate buffer will enhance staining.

Likely to work with frozen sections.

In some cases, the antibody may be diluted further than indicated.

IHC HUMAN CONTROLS Breast Carcinoma, Colon Carcinoma, Laryngeal Squamous Cell Carcinoma, Skin, Small Cell Lung

Cancer, Testicular Seminoma


IHC MOUSE CONTROLS Squamous Cell Carcinoma

ADDITIONAL INFO https://www.bethyl.com/product/IHC-00014-T


Use the link above to view SDS, a current list of citations, and other product specific information.

This document certifies that this product has met all of the quality control standards defined by Bethyl Laboratories, Inc. Eric McIntush, PhD | Chief Scientific Officer Date: June 21, 2019







Detection of human phospho–MCM2 (Ser 108) by immunohistochemistry. Sample: FFPE section of human seminoma. Antibody: Affinity purified rabbit anti–Phospho–MCM2 (Ser 108) (Cat. No. IHC–00014–T) used at a dilution of 1:100. Detection: DAB



Detection of mouse phospho–MCM2 (Ser 108) by immunohistochemistry. *Sample:* FFPE section of mouse squamous cell carcinoma. *Antibody:* Affinity purified rabbit anti–Phospho–MCM2 (Ser 108) (Cat. No. IHC–00014–T) used at a dilution of 1:100. *Detection:* DAB



Detection of human Phospho MCM2 (Ser 108) by immunohistochemistry. Sample: FFPE section of human colon carcinoma. Antibody: Affinity purified rabbit anti-Phospho-MCM2 (Ser 108) (Cat. No. IHC-00014-T) used at a dilution of 1:100. Detection: Red-fluorescent Alexa Fluor® 555 goat antirabbit IgG (Invitrogen) used at a dilution of 1:500.



Detection of human Phospho MCM2 (Ser 108) by immunohistochemistry. *Sample:* FFPE section of human small cell lung cancer. *Antibody:* Affinity purified rabbit anti-Phospho-MCM2 (Ser 108) (Cat. No. IHC-00014-T) used at a dilution of 1:100. *Detection:* Red-fluorescent Alexa Fluor® 555 goat antirabbit IgG (Invitrogen) used at a dilution of 1:500.